

The epigenetics of normal pregnancy

Jonathan D Best PhD and **Nessa Carey** PhD

CellCentric, Chesterford Research Park, Little Chesterford CB10 1XL, UK

Summary: Epigenetic modifications to chromatin are essential for the specification and maintenance of cell fate, enabling the same genome to programme a variety of cellular outcomes. Epigenetic modulation of gene expression is also a critical mechanism by which cells stabilize their responses to environmental stimuli, including both nutritional cues and hormonal signalling. Unsurprisingly, epigenetics is proving to be vitally important in fetal development, and this review addresses our current understanding of the roles of epigenetic regulation in the prenatal phase. It is striking that while there has been a major interest in the intersection of fetal health with epigenetics, there has been relatively little discussion in the literature on epigenetic changes in the pregnant woman, and we attempt to redress this balance, drawing on the fragmented but intriguing experimental literature in this field.

Keywords: endocrinology, metabolism, nutrition, physiology

INTRODUCTION TO EPIGENETICS

Epigenetic traits have been defined operationally as 'stably heritable phenotypes resulting from changes in a chromosome without alterations in the DNA sequence'.¹ Epigenetic modifications (also known as epigenetic marks) form a network of covalent alterations to DNA and histone proteins (Figure 1) which, in turn, interacts with other cellular proteins, typically in multicomponent mediator complexes. The end result is regulation of gene expression. There are a number of excellent recent reviews of the molecular biology of epigenetic control of gene expression.^{2,3}

Developmental biology is arguably the discipline that has had the greatest impact on spotlighting the fundamental nature of epigenetic control of cell fate. From the amphibian somatic cell nuclear transfer experiments of John Gurdon starting in the 1960s, through Azim Surani's ground-breaking studies on the non-equivalence of the maternal and paternal genomes, right up to the current excitement following Shinya Yamanaka's creation of induced pluripotent stem cells, developmental biology has been at the forefront of the great conceptual breakthroughs in epigenetics.

The regulation of gene expression via epigenetic modifications may be relatively short-term and dynamic, or may be exceptionally stable if the chromatin modifications lead to the hypermethylated DNA state associated with the formation of transcriptionally silent heterochromatin. It is clear that epigenetics plays a major role in embryonic development. The transient modifications function in response to signalling molecules to convert relatively undifferentiated cells into increasingly specialized moieties. The gene expression patterns can be stabilized by the development of differential DNA methylation of specific genes, locking cells into a particular fate. These processes are absolutely essential in the generation

of a new multitrillion celled human from a single-celled pluripotent zygote.⁴

Pregnancy is guided by a well-known interaction of hormonal changes which not only ensures that key features of the developmental process (such as placentation) occur at the right time but also that parturition occurs once the baby's development is complete. Acting in concert with the underlying hormonal aspects of pregnancy is the complex interplay of epigenetic modifications, whose appropriate integration and critical timing are also vital for a successful pregnancy.

Molecular epigenetics has struggled to elucidate the key mediators of all these stages. When we consider the size and complexity of the human genome and epigenome, in which there are multiple possible combinations of epigenetic modifications at any given chromatin locus, the reasons for this become obvious. However, recent technological advances are generating enormous amounts of data that are opening windows into the control of fetal development. There has been much less attention paid to the effects of pregnancy on the mother, even though epigenetic modifications to chromatin are clearly an important component of long-term responses to hormonal signalling.


In this review, we attempt to provide an overview of the recent advances in our understanding of the molecular epigenetic regulation of pregnancy, with respect to both the fetal and maternal perspective. We do not address the topic of imprinting, as a number of excellent reviews already exist for this aspect of developmental epigenetics.^{5,6} Our focus in this review is on healthy pregnancy. The role of epigenetic processes in abnormal pregnancy will be the subject of a separate review.

EPIGENETIC INFLUENCES ON FETAL DEVELOPMENT

Ovulation

Clearly, ovulation does not always result in pregnancy, but the former is a prerequisite for the latter, so it is instructive to

Correspondence to: Dr Jonathan D Best
Email: jonathan.best@cellcentric.com



consider the epigenetics of this precursor to the pregnant state. The first stage of pregnancy requires the production of a viable oocyte from the ovaries, its fusion with a sperm cell and implantation into the wall of the uterus. Oocyte production begins in the primordial germ cells and requires erasure of the majority of established somatic epigenomic modifications.³ In mice, the histone methyltransferase (HMT) MLL2 is required for female fertility and development of oocytes particularly postnatally. MLL2 generates trimethylation of lysine position 4 on histone H3 (H3K4me3), a modification that is associated with activation of gene expression. The MLL2 protein is expressed at high levels in the developing oocytes.⁷ Its loss via a conditional knockout results in female mice which fail to ovulate. Maternally contributed expression of epigenetic enzymes is not restricted to MLL2. For example, the protein arginine methyltransferase, PRMT5, which generates repressive histone modifications, is also supplied by the female gamete.⁸ It is likely that such enzymes play key roles at very early points in development. This preloading with epigenetic enzymes is important because, once formed, the oocytes are transcriptionally silenced until fertilization.

As ovulation approaches, a number of epigenetic changes take place during the growth of oocytes with increases detected in DNA methylation along with a number of histone lysine acetylation and methylation events on histones H3 and H4.⁹ These changes correlate with the presence of a number of histone acetyltransferase enzymes including Src-1, p300 and P-caf. The changes in histone methylation suggest possible roles for HMTs, Smyd3, Set7 and Mll2 proteins. However, there is not always a simple relationship between the level of expression of HMTs and the pattern of methylation at their

histone targets. This has been demonstrated for the methylation of H3K9. G9a and Glp, which dimethylate H3K9, are present at high levels along with elevated levels of H3K9me2. However, Eset and Suv39h, which are responsible for trimethylation, are also present at high levels but levels of H3K9me3 are low. This suggests that specific roles of these enzymes are not purely driven by their levels of expression.

Preimplantation

After release of the mature oocyte and fertilization by the sperm cell, the newly formed zygote undergoes a number of developmental changes which results in the epigenetic reprogramming of the non-imprinted genes in the genome. This re-sets the epigenome to an expression status compatible with totipotency (the ability of a cell to form all cell types in the body) before implantation (reviewed in reference¹⁰). One of the key mechanisms, but which is least understood, is the initial large-scale erasure of DNA methylation marks from the paternal genome. This would appear to be an active process as it occurs before replication and cell division takes place. However the mechanism responsible for this (whether direct DNA demethylation or indirect changes via base excision response) remains controversial (fully detailed in reference³). Recent work in embryonic stem (ES) cells demonstrates this is most likely to involve hydroxylation of the methylated cytosine residue, mediated by the Tet family of enzymes.¹¹

Following this epigenomic erasure, the epigenome is re-patterned with *de novo* establishment of histone and DNA modifications as cell lineages develop. This occurs from the

very earliest stages of cellular commitment, during formation of the blastocyst. This is functionally subdivided into the outer layer of differentiated cells (the trophoectoderm [TE]), surrounding the small cluster of cells in the inner cell mass (ICM) which contains the pluripotent cells. The ICM is the experimental source of ES cells. Some of the epigenetic mechanisms underlying this process are beginning to be revealed (reviewed in detail in reference¹²).

Much of the work studying this reprogramming has been conducted in rodents where the differentiation of these early cells into the TE and the ICM is correlated with the expression of *Cdx2* and *Oct4*, respectively. The downstream effects of *Oct4* activity are critically dependent on the establishment of repressive histone modifications by the H3K9 HMT Eset (also known as *Setdb1*). Eset is recruited by and co-associates with *Oct4* to repress *Cdx2* via methylation of H3K9. Eset depletion causes embryonic lethality at the peri-implantation stage. This is associated with incomplete formation of the ICM, with the cells becoming more TE-like due to the de-repression of *Cdx2*.^{13–15}

Extreme early embryonic lethality in response to disruption of key epigenetic mediators is not restricted to Eset. Knockout of *Prmt5* also results in exceptionally early loss of embryos.⁸ For each of these genes, disruption of the development of the ICM is so severe that it is impossible to generate ES cells from the homozygous knockouts.

Timing of the action of these and other epigenetic enzymes is critical to normal embryo development. However, the precise roles of these various epigenetic agents in initiating, driving or maintaining developmental pathways remain the subject of intense scrutiny.¹⁶

Implantation and placentation

The next stage for the developing zygote is its implantation into the endometrial wall and the formation of the placenta. As described above, formation of the trophoblast cells is critically dependent on the up-regulation of *Cdx2* (reviewed in reference¹⁷). One of the factors that influence *Cdx2* expression is *Elf5*. In trophoblast cells, the promoter region of the *Elf5* gene is hypomethylated, resulting in increased levels of *Elf5* which drive up expression of *Cdx2*. This results in a positive feedback loop between *Cdx2* and *Elf5* expression where *Elf5* drives *Cdx2*, and where *Cdx2* in turn drives *Elf5*, resulting in trophoblast differentiation. In contrast, in the cells of the ICM, the DNA of the *Elf5* promoter is hypermethylated, which shuts down *Elf5* transcription, thereby preventing trophoblast formation.¹⁸

Unlike the cells that arise from the ICM, the genome of the placental precursors remains relatively free from *de novo* DNA methylation (reviewed in reference¹⁹). The key factors for causing DNA methylation of those genes in the placenta which do become methylated are still being elucidated,^{20–22} but interestingly distinct changes are seen in histone methylation marks in the different cell lineages of the developing zygote.²³ Preimplantation, the repressive H3K27me3 mark, is present at relatively low levels in the TE and extra-ES stem cells compared with the ICM-derived ES cells. This is consistent with the relatively low levels of the enzymes that constitute the relevant methyltransferase machinery, the polycomb repressive complex 2. Postimplantation, the levels of H3K27me3 are relatively high in extraembryonic tissues. Together with the presence of other repressive marks in these tissues, such as

H3K9me3, this suggests that these modifications are associated with the formation of specific cell lineages.²³

Nutrition as an epigenetic stimulus in fetal development

It is during the first trimester that the majority of the fetal development occurs and this coincides with global changes to epigenetic modifications in the embryo.³ How these widespread changes relate to specific pathways of fetal tissue development remains a key question. One productive area of research has been the differentiation of mesenchymal cells in chondrogenesis and osteogenesis.^{24,25} These studies have revealed the importance of histone acetylation and DNA methylation modifications in mediating cellular differentiation.

Another area of great interest in relation to pregnancy is the impact of the environment on epigenetic control. For example, the formation and closure of the neural tube is a very important event in the embryo's development, and incomplete development can lead to malformations referred to as neural tube defects.²⁶ Deficiencies in the intake of folic acid have long been linked to neural tube defects. Intriguingly, recent studies have suggested this may have an epigenetic component (reviewed in reference²⁷). Folic acid is required for the production of *S*-adenosyl methionine (SAM). SAM is the molecule that donates the methyl group when DNA methyltransferases modify DNA (see Figure 1a). Mice weaned on a diet low in folic acid develop abnormal regulation of imprinted regions of the genome,²⁸ demonstrating that the levels of this key nutrient do influence epigenetic signalling.

The effects of the nutritional status of the mother on fetal outcomes have been intensively studied, particularly since the landmark Dutch famine study.²⁹ This revealed significant effects on resulting adult obesity in the offspring (reviewed in reference³⁰). These effects have been linked to DNA methylation events on a number of growth-related genes and recently to placental expression of micro-RNAs.³¹ Similar to folic acid, decreases in maternal dietary choline intake negatively influence the availability of SAM as a DNA methyl donor, resulting in epigenetic changes to the offspring.³²

EPIGENETIC CHANGES TO THE PREGNANT MOTHER

Oestrogen signalling

As the ovarian follicles mature, they secrete increasing amounts of oestrogens, and it is well documented that these hormones have a number of effects on the pregnant female.³³ Oestradiol, the predominant oestrogen, acts to enhance the effects of follicle stimulating hormone. This promotes follicle maturation and development of breast tissue, and initiates the proliferation of cells in the formation of the endometrium.^{34–36} Much of the information regarding epigenetic interaction with oestrogen signalling has been derived from cancer cells lines.^{37–40} However, the fact that both processes are strongly associated with the proliferation of cell numbers, albeit with entirely different endpoints, may allow us to draw parallels between these systems and give us some additional insights into the epigenetic regulation in response to the types of hormonal changes seen during pregnancy.

The effects of oestrogens are mediated by the alpha and beta oestrogen receptors. Following ligand binding, these translocate from the cell membrane to the nucleus, where they bind directly or as part of a multimeric complex to DNA. This generally promotes gene expression, but the effects can be further modulated by interaction with a number of epigenetic enzymes.^{41,42} This results in the establishment of multimeric protein complexes, targeted to specific genomic locations, which leads to an epigenetic signalling cascade.

The interaction of epigenetic processes with hormonal signalling can be more complicated than a simple analysis of gene function might indicate. For example, KMT8 (as known as PRDM2 or RIZ1) methylates H3K9, generating a repressive chromatin mark. Unexpectedly, KMT8 acts as a co-activator with the oestrogen receptor. KMT8 knockout in female mice results in a number of phenotypes consistent with a decreased response to female sex hormones. In pregnant females these include a reduced increase in uterine weight, reduced epithelial vaginal thickening in response to oestrogen and reduced litter sizes compared with their wild-type counterparts.⁴³

More recently, modifications generated by other HMTs have been shown to be important in the oestrogen signalling pathways. The arginine methyltransferases PRMT1, PRMT4 and PRMT6 which methylate histone proteins are each co-activators of receptor-mediated oestrogen signalling in breast-derived cell lines.^{37,40} Among the lysine methyltransferases, SMYD3 and SET7 play co-activator roles with the oestrogen receptor alpha, when exposed to oestrogen. SMYD3 is responsible for H3K4 methylation, generating an activating chromatin modification and thereby facilitating oestrogen-mediated gene activation at ER-induced target genes.³⁸ SET7 may influence oestrogen signalling via a different mechanism. It has been reported that SET7 directly methylates the oestrogen receptor itself at lysine 302. This stabilizes the receptor, promoting its efficient recruitment to target genes.³⁹ This highlights an important additional layer of complexity, which must be considered when analysing the effects and roles of epigenetic enzymes. Many of these proteins have both histone and non-histone targets, and unravelling which of these modes of action is most critical in response to a specific stimulus can be challenging.

Successful implantation is critical for the continuing effective development of the embryo and adequate uterine receptivity plays a major role. The formation of the endometrium is important to this successful outcome and is linked to changes in the acetylation of histone lysine residues which occur during the menstrual cycle. Interestingly, the histone deacetylase inhibitor sodium valproate decreases the proliferation of an endometrial cell line, potentially linking this histone modification to endometrial formation (all above reviewed in reference³⁴). The decidualization of the endometrial stromal cells which regulates the invasion of the trophoblast cells has been reported to be in part regulated by DNA methylation,^{44,45} although any further role of DNA methylation in endometrial function has yet to be fully characterized.⁴⁶

As pregnancy progresses, a number of overt changes become increasingly obvious in the mother. The breast tissue changes in preparation for lactation, a process that is largely underpinned by lactogenic hormones secreted via the anterior pituitary. However, there is evidence that many of these changes are mediated at least in part by epigenetic mechanisms. The promoters of the milk-related genes have low levels of DNA methylation in lactating mammary glands, allowing gene expression

to take place. In non-mammary or non-lactating mammary tissue, these genes are hypermethylated at the DNA level (for a review see reference⁴⁷). In the mammary glands of mice, levels of Dnmt1 and Dnmt3b3b are lower in lactating females compared with virgin and involuted females.⁴⁸

Lactation is also linked to histone acetylation. The casein gene cluster shows high levels of histone acetylation in the mammary tissue of lactating mice. Such histone acetylation is typically associated with transcriptionally active genes.⁴⁷ Other studies have shown changes in nuclear architecture during lactation. H4K20me3 foci become more compact in lactating mammary epithelial cells and located more frequently at the nuclear periphery. In contrast, H4K20me3 foci are found throughout the nucleus in mammary epithelial cells from virgin and pregnant mice.⁴⁹

CONCLUSIONS

It is clear that epigenetic regulation is a key feature of pregnancy and development, but our current understanding is rather piecemeal. In addition to the histone and DNA modifications that are described in this review, there are further aspects which are currently underexplored. These include the roles of non-canonical histone variants and the impact of regulatory non-coding RNAs, to name but two. In addition, most of the fundamental insights into the roles of epigenetics have been obtained from rodent species. While it is ethically and scientifically challenging to research reproduction in humans, it is essential that we gain a better mechanistic understanding of how these processes operate in human reproduction. This will be key for the improvement of both infant and maternal health, particularly in a more rational analysis of dietary effects in pregnancy. Furthermore, the only currently marketed drugs targeting epigenetic proteins are licensed in oncology. This is likely to change as epigenetic drug discovery progresses in chronic conditions. If such drugs are to be administered during pregnancy or to women of childbearing age, they will create a greater need for enhanced mechanistic understanding of epigenetic regulation of development and pregnancy.

DECLARATIONS

Competing interests: JB is an employee of CellCentric Ltd and NC is an employee of Pfizer Ltd.

Funding: N/A.

Ethical approval: N/A.

Guarantor: JB.

Contributorship: JB and NC co-wrote the manuscript. Both authors reviewed and edited the manuscript and approved the final version of the manuscript.

Acknowledgements: N/A.

REFERENCES

- Berger SL, Kouzarides T, Shiekhattar R, Shilatifard A. An operational definition of epigenetics. *Genes Dev* 2009;23:781–3. Epub 2009/04/03
- Cheng X, Blumenthal RM. Coordinated chromatin control: structural and functional linkage of DNA and histone methylation. *Biochemistry* 2010;49:2999–3008. Epub 2010/03/10
- Feng S, Jacobsen SE, Reik W. Epigenetic reprogramming in plant and animal development. *Science* 2010;330:622–7. Epub 2010/10/30

- 4 Ikegami K, Ohgane J, Tanaka S, Yagi S, Shiota K. Interplay between DNA methylation, histone modification and chromatin remodeling in stem cells and during development. *Int J Dev Biol* 2009;**53**:203–14. Epub 2009/05/05
- 5 Frost JM, Moore GE. The importance of imprinting in the human placenta. *PLoS Genet* 2010;**6**:e1001015. Epub 2010/07/10
- 6 McEwen KR, Ferguson-Smith AC. Distinguishing epigenetic marks of developmental and imprinting regulation. *Epigenet Chromatin* 2010;**3**:2. Epub 2010/02/26
- 7 Andreu-Vieyra CV, Chen R, Agno JE, *et al.* MLL2 is required in oocytes for bulk histone 3 lysine 4 trimethylation and transcriptional silencing. *PLoS Biol* 2010;**8**. Epub 2010/09/03
- 8 Tee WW, Pardo M, Theunissen TW, *et al.* Prmt5 is essential for early mouse development and acts in the cytoplasm to maintain ES cell pluripotency. *Genes Dev* 2010;**24**:2772–7. Epub 2010/12/17
- 9 Kageyama S, Liu H, Kaneko N, Oga M, Nagata M, Aoki F. Alterations in epigenetic modifications during oocyte growth in mice. *Reproduction* 2007;**133**:85–94. Epub 2007/01/25
- 10 Burton A, Torres-Padilla ME. Epigenetic reprogramming and development: a unique heterochromatin organization in the preimplantation mouse embryo. *Brief Funct Genomics* 2010;**9**:444–54. Epub 2010/12/28
- 11 Ficiz G, Branco MR, Seisenberger S, *et al.* Dynamic regulation of 5-hydroxymethylcytosine in mouse ES cells and during differentiation. *Nature* 2011. Epub 2011/04/05
- 12 Duranthon V, Watson AJ, Lonergan P. Preimplantation embryo programming: transcription, epigenetics, and culture environment. *Reproduction* 2008;**135**:141–50. Epub 2008/02/02
- 13 Lohmann F, Loureiro J, Su H, *et al.* KMT1E mediated H3K9 methylation is required for the maintenance of embryonic stem cells by repressing trophoblast differentiation. *Stem Cells* 2010;**28**:201–12. Epub 2009/12/17
- 14 Yeap LS, Hayashi K, Surani MA. ERG-associated protein with SET domain (ESET)-Oct4 interaction regulates pluripotency and represses the trophoblast lineage. *Epigenet Chromatin* 2009;**2**:12. Epub 2009/10/09
- 15 Yuan P, Han J, Guo G, *et al.* Eset partners with Oct4 to restrict extraembryonic trophoblast lineage potential in embryonic stem cells. *Genes Dev* 2009;**23**:2507–20. Epub 2009/11/04
- 16 Combes AN, Whitelaw E. Epigenetic reprogramming: enforcer or enabler of developmental fate? *Dev Growth Differ* 2010;**52**:483–91. Epub 2010/07/09
- 17 Hemberger M. Genetic-epigenetic intersection in trophoblast differentiation: implications for extraembryonic tissue function. *Epigenetics* 2010;**5**:24–9. Epub 2010/01/20
- 18 Ng RK, Dean W, Dawson C, *et al.* Epigenetic restriction of embryonic cell lineage fate by methylation of Elf5. *Nat Cell Biol* 2008;**10**:1280–90. Epub 2008/10/07
- 19 Nelissen EC, van Montfoort AP, Dumoulin JC, Evers JL. Epigenetics and the placenta. *Hum Reprod Update* 2010. Epub 2010/10/21
- 20 Grigoriu A, Ferreira JC, Choufani S, Baczyk D, Kingdom J, Weksberg R. Cell specific patterns of methylation in the human placenta. *Epigenetics* 2011;**6**:53–64. Epub 2010/12/07
- 21 Maccani MA, Marsit CJ. Epigenetics in the placenta. *Am J Reprod Immunol* 2009;**62**:78–89. Epub 2009/07/21
- 22 Novakovic B, Wong NC, Sibson M, *et al.* DNA methylation-mediated down-regulation of DNA methyltransferase-1 (DNMT1) is coincident with, but not essential for, global hypomethylation in human placenta. *J Biol Chem* 2010;**285**:9583–93. Epub 2010/01/15
- 23 Rugg-Gunn PJ, Cox BJ, Ralston A, Rossant J. Distinct histone modifications in stem cell lines and tissue lineages from the early mouse embryo. *Proc Natl Acad Sci USA* 2010;**107**:10783–90. Epub 2010/05/19
- 24 Furumatsu T, Ozaki T. Epigenetic regulation in chondrogenesis. *Acta Med Okayama* 2010;**64**:155–61. Epub 2010/07/03
- 25 Hsiao SH, Lee KD, Hsu CC, *et al.* DNA methylation of the Trip10 promoter accelerates mesenchymal stem cell lineage determination. *Biochem Biophys Res Commun* 2010;**400**:305–12. Epub 2010/08/24
- 26 Dhaulakhandi DB, Rohilla S, Rattan KN. Neural tube defects: review of experimental evidence on stem cell therapy and newer treatment options. *Fetal Diagn Ther* 2010;**28**:72–8. Epub 2010/08/07
- 27 Blom HJ. Folic acid, methylation and neural tube closure in humans. *Birth Defects Res A Clin Mol Teratol* 2009;**85**:295–302. Epub 2009/03/21
- 28 Waterland RA, Lin JR, Smith CA, Jirtle RL. Post-weaning diet affects genomic imprinting at the insulin-like growth factor 2 (Igf2) locus. *Hum Mol Genet* 2006;**15**:705–16. Epub 2006/01/20
- 29 Ravelli GP, Stein ZA, Susser MW. Obesity in young men after famine exposure in utero and early infancy. *N Engl J Med* 1976;**295**:349–53. Epub 1976/08/12
- 30 Simmons R. Epigenetics and maternal nutrition: nature v. nurture. *Proc Nutr Soc* 2011;**70**:73–81. Epub 2010/11/30
- 31 Maccani MA, Padbury JF, Marsit CJ. miR-16 and miR-21 expression in the placenta is associated with fetal growth. *PLoS One* 2011;**6**:e21210. Epub 2011/06/24
- 32 Zeisel SH. Epigenetic mechanisms for nutrition determinants of later health outcomes. *Am J Clin Nutr* 2009;**89**:1488S–93S. Epub 2009/03/06
- 33 Albrecht ED, Pepe GJ. Estrogen regulation of placental angiogenesis and fetal ovarian development during primate pregnancy. *Int J Dev Biol* 2010;**54**:397–408. Epub 2009/10/31
- 34 Munro SK, Farquhar CM, Mitchell MD, Ponnampalam AP. Epigenetic regulation of endometrium during the menstrual cycle. *Mol Hum Reprod* 2010;**16**:297–310. Epub 2010/02/09
- 35 Richards JS, Pangas SA. The ovary: basic biology and clinical implications. *J Clin Invest* 2010;**120**:963–72. Epub 2010/04/07
- 36 Tamm K, Room M, Salumets A, Metsis M. Genes targeted by the estrogen and progesterone receptors in the human endometrial cell lines HEC1A and RL95–2. *Reprod Biol Endocrinol* 2009;**7**:150. Epub 2009/12/26
- 37 Harrison MJ, Tang YH, Dowhan DH. Protein arginine methyltransferase 6 regulates multiple aspects of gene expression. *Nucleic Acids Res* 2010;**38**:2201–16. Epub 2010/01/06
- 38 Kim H, Heo K, Kim JH, Kim K, Choi J, An W. Requirement of histone methyltransferase SMYD3 for estrogen receptor-mediated transcription. *J Biol Chem* 2009;**284**:19867–77. Epub 2009/06/11
- 39 Subramanian K, Jia D, Kapoor-Vazirani P, *et al.* Regulation of estrogen receptor alpha by the SET7 lysine methyltransferase. *Mol Cell* 2008;**30**:336–47. Epub 2008/05/13
- 40 Wagner S, Weber S, Kleinschmidt MA, Nagata K, Bauer UM. SET-mediated promoter hypoacetylation is a prerequisite for coactivation of the estrogen-responsive pS2 gene by PRMT1. *J Biol Chem* 2006;**281**:27242–50. Epub 2006/07/25
- 41 Leader JE, Wang C, Popov VM, Fu M, Pestell RG. Epigenetics and the estrogen receptor. *Ann NY Acad Sci* 2006;**1089**:73–87. Epub 2007/01/31
- 42 Yap OW, Bhat G, Liu L, Tollefsbol TO. Epigenetic modifications of the Estrogen receptor beta gene in epithelial ovarian cancer cells. *Anticancer Res* 2009;**29**:139–44. Epub 2009/04/01
- 43 Carling T, Kim KC, Yang XH, Gu J, Zhang XK, Huang S. A histone methyltransferase is required for maximal response to female sex hormones. *Mol Cell Biol* 2004;**24**:7032–42. Epub 2004/07/30
- 44 Logan PC, Ponnampalam AP, Rahnama F, Lobie PE, Mitchell MD. The effect of DNA methylation inhibitor 5-Aza-2'-deoxycytidine on human endometrial stromal cells. *Hum Reprod* 2010;**25**:2859–69. Epub 2010/09/09
- 45 Vincent ZL, Farquhar CM, Mitchell MD, Ponnampalam AP. Expression and regulation of DNA methyltransferases in human endometrium. *Fertil Steril* 2010. Epub 2010/10/26
- 46 Furst RW, Meyer HH, Schweizer G, Ulbrich SE. Is DNA methylation an epigenetic contribution to transcriptional regulation of the bovine endometrium during the estrous cycle and early pregnancy? *Mol Cell Endocrinol* 2011. Epub 2011/08/02
- 47 Rijnkels M, Kabotyanski E, Montazer-Torbati MB, *et al.* The epigenetic landscape of mammary gland development and functional differentiation. *J Mammary Gland Biol Neoplasia* 2010;**15**:85–100. Epub 2010/02/17
- 48 Tsellou E, Michailidi C, Pafiti A, Troungos C. DNA methylation-independent regulation of p16 in epithelial cells during mouse mammary gland development. *Epigenetics* 2008;**3**:143–8. Epub 2008/06/24
- 49 Kress C, Ballester M, Devinoy E, Rijnkels M. Epigenetic modifications in 3D: nuclear organization of the differentiating mammary epithelial cell. *J Mammary Gland Biol Neoplasia* 2010;**15**:73–83. Epub 2010/02/10

(Accepted 7 September 2011)